

POLICY Document for COSENTYX (Secukinumab)

The overall objective of this policy is to support the appropriate and cost-effective use of the medication, specific to use of preferred medication options, and overall, clinically appropriate use. This document provides specific information to both sections of the overall policy.

Section 1: Preferred Product

- Policy information specific to preferred medications

Section 2: Clinical Criteria

- Policy information specific to the clinical appropriateness for the medication

Section 1: Preferred Product

CAREFIRST: EXCEPTIONS CRITERIA AUTOIMMUNE CONDITIONS

PRIMARY PREFERRED PRODUCTS: ENTYVIO, SIMPONI ARIA, SKYRIZI, STELARA

Client Requested: The intent of the criteria is to ensure that patients follow selection elements as established by CareFirst.

POLICY

This policy informs prescribers of preferred products and provides an exception process for targeted products through prior authorization.

I. PLAN DESIGN SUMMARY

This program applies to the autoimmune drug products specified in this policy. Coverage for targeted product is provided based on clinical circumstances that would exclude the use of the preferred products and may be based on previous use of a product. The coverage review process will ascertain situations where a clinical exception can be made. This program applies to all members requesting treatment with a targeted product.

Each referral is reviewed based on all utilization management (UM) programs implemented for the client.

Table. Autoimmune Products

	Product(s)
Preferred	<ul style="list-style-type: none"> • Entyvio (vedolizumab) • Simponi Aria (golimumab, intravenous) • Skyrizi (risankizumab-rzaa) • Stelara (ustekinumab)
Targeted	<ul style="list-style-type: none"> • Actemra (tocilizumab) • Cimzia (certolizumab pegol) • Cosentyx (Secukinumab) • Ilumya (tildrakizumab-asmn)

	<ul style="list-style-type: none"> • Orencia (abatacept) • Tofidence (Tocilizumab-bavi) • Tremfya (guselkumab) • Tyenne (Tocilizumab-aazg) • Tysabri (natalizumab)
--	--

*: Medications considered formulary or preferred on your plan may still require a clinical prior authorization review

II. EXCEPTION CRITERIA

This program applies to members requesting treatment for an indication that is FDA-approved for the preferred products.

Coverage for a targeted product is provided when any of the following criteria is met:

- A. For Actemra, Tofidence, or Tyenne when any of the following criteria is met:
 1. When the request is for Systemic Juvenile Idiopathic Arthritis
 2. When the request is for Giant Cell Arteritis
 3. Member is currently receiving treatment with the requested targeted product, excluding when the requested targeted product is obtained as samples or via manufacturer's patient assistance programs.
 4. Member has a documented inadequate response, contraindication, or intolerable adverse event to Simponi Aria.
- B. For Cimzia, when any of the following criteria is met:
 1. When the request is for Axial Spondylarthritis
 2. Member is pregnant, breastfeeding, or of childbearing potential
 3. Member suffers from Trypanophobia (needle-phobic) and cannot self-inject
 4. Member is currently receiving treatment with the requested targeted product, excluding when the requested targeted product is obtained as samples or via manufacturer's patient assistance programs.
 5. Member has a documented inadequate response, contraindication, or intolerable adverse event to Simponi Aria, Stelara, Entyvio, and Skyrizi.
- C. For Cosentyx, when any of the following criteria is met:
 1. When the request is for Axial Spondylarthritis
 2. Member is currently receiving treatment with the requested targeted product, excluding when the requested targeted product is obtained as samples or via manufacturer's patient assistance programs.
 3. Member has a documented inadequate response, contraindication, or intolerable adverse event to Simponi Aria, Stelara, and Skyrizi.
- D. For Ilumya, when any of the following criteria is met:
 1. Member is currently receiving treatment with the requested targeted product, excluding when the requested targeted product is obtained as samples or via manufacturer's patient assistance programs.
 2. Member has a documented inadequate response, contraindication, or intolerable adverse event to Stelara and Skyrizi
- E. For Orencia, when any of the following criteria is met:
 1. Member is currently receiving treatment with the requested targeted product, excluding when the requested targeted product is obtained as samples or via manufacturer's patient assistance programs.
 2. Member has a documented inadequate response, contraindication, or intolerable adverse event to Simponi Aria, Stelara, and Skyrizi.
- F. For Tremfya, when any of the following criteria is met:
 1. Member is currently receiving treatment with the requested targeted product, excluding when the requested targeted product is obtained as samples or via manufacturer's patient assistance programs.

2. Member has a documented inadequate response, contraindication, or intolerable adverse event to Simponi Aria, Stelara, Entyvio, and Skyrizi.

G. For Tysabri, when any of the following criteria is met:

1. When the request is for Multiple Sclerosis
2. Member is currently receiving treatment with the requested targeted product, excluding when the requested targeted product is obtained as samples or via manufacturer's patient assistance programs.
3. Member has a documented inadequate response, contraindication, or intolerable adverse event to Stelara, Entyvio, and Skyrizi.

Section 2: Clinical Criteria

Specialty Guideline Management

Cosentyx

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Cosentyx	secukinumab

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications¹

- Moderate to severe plaque psoriasis (PsO) in patients 6 years of age and older who are candidates for systemic therapy or phototherapy
- Active psoriatic arthritis (PsA) in patients 2 years of age and older
- Adults with active ankylosing spondylitis (AS)
- Adults with active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation
- Active enthesitis-related arthritis (ERA) in patients 4 years of age and older
- Adults with moderate to severe hidradenitis suppurativa (HS)

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

Plaque psoriasis (PsO)

Initial requests

Chart notes or medical record documentation of affected area(s) and body surface area (BSA) affected (if applicable).

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Continuation requests

Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.

Psoriatic arthritis (PsA), ankylosing spondylitis (AS), non-radiographic axial spondyloarthritis (nr-axSpA), enthesitis-related arthritis (ERA), and hidradenitis suppurativa

Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

Prescriber Specialties

This medication must be prescribed by or in consultation with one of the following:

Plaque psoriasis: dermatologist

Psoriatic arthritis and hidradenitis suppurativa: rheumatologist or dermatologist

Ankylosing spondylitis, non-radiographic axial spondyloarthritis, and enthesitis-related arthritis: rheumatologist

Coverage Criteria

Plaque psoriasis (PsO)^{1-2,8,13,14}

Authorization of 12 months may be granted for members 6 years of age or older who have previously received a biologic or targeted synthetic drug (e.g., Sotyktu, Otezla) indicated for the treatment of moderate to severe plaque psoriasis.

Authorization of 12 months may be granted for members 6 years of age or older for the treatment of moderate to severe plaque psoriasis when any of the following criteria is met:

- Crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
- At least 10% of body surface area (BSA) is affected.
- At least 3% of body surface area (BSA) is affected and the member meets any of the following criteria:
 - Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin.
 - Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine, and acitretin (see Appendix).

Psoriatic arthritis (PsA)^{1,3,4,10,13}

Authorization of 12 months may be granted for members 2 years of age or older who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Otezla) indicated for active psoriatic arthritis.

Authorization of 12 months may be granted for members 2 years of age or older for treatment of active psoriatic arthritis when either of the following criteria is met:

- Member has mild to moderate disease and meets one of the following criteria:
 - Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.
 - Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix), or another conventional synthetic drug (e.g., sulfasalazine).
 - Member has enthesitis or predominantly axial disease.
- Member has severe disease.

Ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA)^{1,5-7,15}

Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Xeljanz) indicated for active ankylosing spondylitis or active non-radiographic axial spondyloarthritis.

Authorization of 12 months may be granted for adult members for treatment of active ankylosing spondylitis or active non-radiographic axial spondyloarthritis when any of the following criteria is met:

- Member has had an inadequate response to at least two nonsteroidal anti-inflammatory drugs (NSAIDs).
- Member has an intolerance or contraindication to two or more NSAIDs.

Enthesitis-related arthritis (ERA)^{1,11,12}

Authorization of 12 months may be granted for members 4 years of age or older who have previously received a biologic for treatment of active enthesitis-related arthritis.

Authorization of 12 months may be granted for members 4 years of age or older for treatment of active enthesitis-related arthritis when both of the following criteria are met:

- Member has active disease demonstrated by at least three active joints involved and at least one site of active enthesitis at baseline or documented by history.
- Member meets either of the following:
 - Member has had an inadequate response to nonsteroidal anti-inflammatory drugs (NSAIDs), sulfasalazine, or methotrexate.
 - Member has an intolerance or contraindication to NSAIDs, sulfasalazine (e.g., porphyria, intestinal or urinary obstruction), and methotrexate (see Appendix).

Hidradenitis suppurativa^{1,16,17}

Authorization of 12 months may be granted for adult members who have previously received a biologic indicated for treatment of moderate to severe hidradenitis suppurativa.

Authorization of 12 months may be granted for adult members for treatment of moderate to severe hidradenitis suppurativa when either of the following is met:

- Member has had an inadequate response to an oral antibiotic used for the treatment of hidradenitis suppurativa for at least 90 days (e.g., clindamycin, metronidazole, moxifloxacin, rifampin, tetracyclines).
- Member has an intolerance or contraindication to oral antibiotics used for the treatment of hidradenitis suppurativa.

Continuation of Therapy

Plaque psoriasis (PsO)¹

Authorization of 12 months may be granted for all members 6 years of age or older (including new members) who are using the requested medication for moderate to severe plaque psoriasis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when either of the following is met:

Reduction in body surface area (BSA) affected from baseline
Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

Psoriatic arthritis (PsA)^{1,10,13}

Authorization of 12 months may be granted for all members 2 years of age or older (including new members) who are using the requested medication for psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Number of swollen joints
- Number of tender joints
- Dactylitis
- Enthesitis
- Axial disease
- Skin and/or nail involvement
- Functional status
- C-reactive protein (CRP)

Ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA)^{1,5-7,15}

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for ankylosing spondylitis or non-radiographic axial spondyloarthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Functional status
- Total spinal pain
- Inflammation (e.g., morning stiffness)
- Swollen joints
- Tender joints
- C-reactive protein (CRP)

Enthesitis-related arthritis (ERA)^{1,11,12}

Authorization of 12 months may be granted for all members 4 years of age or older (including new members) who are using the requested medication for active enthesitis-related arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Number of flares
- Number of joints with active arthritis (e.g., swelling, pain)
- Number of joints with limited movement
- Dactylitis
- Enthesitis

Hidradenitis suppurativa^{1,16,17}

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderate to severe hidradenitis suppurativa and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:

- Reduction in abscess and inflammatory nodule count from baseline
- Reduced formation of new sinus tracts and scarring
- Decrease in frequency of inflammatory lesions from baseline
- Reduction in pain from baseline
- Reduction in suppuration from baseline
- Improvement in frequency of relapses from baseline
- Improvement in quality of life from baseline
- Improvement on a disease severity assessment tool from baseline

Other^{1,9}

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA]) within 12 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

If the screening testing for TB is positive, there must be further testing to confirm there is no active disease (e.g., chest x-ray). Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

Dosage And Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

Appendix

Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine, Acitretin, or Leflunomide¹⁴

- Clinical diagnosis of alcohol use disorder, alcoholic liver disease, or other chronic liver disease
- Drug interaction

Risk of treatment-related toxicity
Pregnancy or currently planning pregnancy
Breastfeeding
Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
Hypersensitivity
History of intolerance or adverse event

REFERENCES:

SECTION 1

1. Actemra [package insert]. South San Francisco, CA: Genentech, Inc.; September 2024.
2. Tofidence [package insert] Cambridge, MA: Biogen MA Inc; July 2024.
3. Tyenne [package insert] Lake Zurich, IL: Fresenius Kabi USA, LLC; September 2024.
4. Cimzia [package insert]. Smyrna, GA: UCB, Inc.; September 2019.
5. Cosentyx [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; August 2024.
6. Entyvio [package insert]. Lexington, MA: Takeda Pharmaceuticals U.S.A., Inc.; April 2024.
7. Ilumya [package insert]. Cranbury, NJ: Sun Pharma Global FZE; April 2024.
8. Orencia [package insert]. Princeton, NJ: Bristol-Meyers Squibb Company; May 2024.
9. Simponi Aria [package insert]. Horsham, PA: Janssen Biotech, Inc.; February 2021.
11. Stelara [package insert]. Horsham, PA: Janssen Biotech, Inc.; March 2024.
12. Tysabri [package insert]. Cambridge, MA: Biogen Inc.; October 2023.
13. Tremfya [package insert]. Horsham, PA; Janssen Biotech, Inc; September 2024
14. Skyrizi [package insert]. North Chicago, IL; AbbVie Inc.; June 2024.

SECTION 2

1. Cosentyx [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; October 2024.
2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. J Am Acad Dermatol. 2011;65(1):137-174.
3. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. Ann Rheum Dis. 2020;79(6):700-712.
4. McInnes IB, Mease PJ, Kirkham B, et al. Secukinumab, a human anti-interleukin-17A monoclonal antibody, in patients with psoriatic arthritis (FUTURE 2): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet. 2015;386(9999):1137-46.
5. Braun J, van den Berg R, Baraliakos, X et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. Ann Rheum Dis. 2011;70:896–904.
6. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. Arthritis Rheumatol. 2019;71(10):1285-1299.
7. Baeten D, Sieper J, Braun J, et al. Secukinumab, an Interleukin-17A Inhibitor, in Ankylosing Spondylitis. N Engl J Med. 2015;373(26):2534-48.
8. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. J Am Acad Dermatol. 2019;80(4):1029-1072.
9. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on January 22, 2025 from: <https://www.cdc.gov/tb/testing/index.html>.
10. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. Arthritis Rheum. 2018;71:5-32.
11. Weiss PF. Diagnosis and treatment of enthesitis-related arthritis. Adolesc Health Med Ther. 2012;2012(3):67-74.

12. Brunner HI, Foeldvari I, Alexeeva E, et al. Secukinumab in enthesitis-related arthritis and juvenile psoriatic arthritis: a randomised, double-blind, placebo-controlled, treatment withdrawal, phase 3 trial. *Ann Rheum Dis*. 2023 Jan;82(1):154-160.
13. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol*. 2022;18(8):465-479.
14. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol*. 2020;82(6):1445-1486.
15. Ramiro S, Nikiphorou E, Sepriano A, et al. ASAS-EULAR recommendations for the management of axial spondyloarthritis: 2022 update. *Ann Rheum Dis*. 2023;82:19-34.
16. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part I: Diagnosis, evaluation, and the use of complementary and procedural management. *J Am Acad Dermatol*. 2019; 81(1): 76-90.
17. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part II: Topical, intralesional, and systemic medical management. *J Am Acad Dermatol*. 2019; 81(1): 91-101.