

POLICY Document for LEQEMBI

The overall objective of this policy is to support the appropriate and cost-effective use of the medication, specific to use of preferred medication options, lower cost site of care and overall, clinically appropriate use. This document provides specific information to each of the three sections of the overall policy.

Section 1: Site of Care

- Policy information specific to site of care (outpatient, hospital outpatient, home infusion)

Section 2: Clinical Criteria

- Policy information specific to the clinical appropriateness for the medication

Section 1: Site of Care

Site of Care Criteria Leqembi

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated.

Brand Name	Generic Name	Dosage Form
Leqembi	lecanemab-irmb	intravenous

Criteria For Approval For Administration In Outpatient Hospital Setting

This policy provides coverage for administration of Leqembi in an outpatient hospital setting for up to 45 days when a member is new to therapy or is reinitiating therapy after not being on therapy for at least 6 months.

This policy provides coverage for administration of Leqembi in an outpatient hospital setting for a longer course of treatment when ANY of the following criteria are met:

- The member has experienced an adverse reaction that did not respond to conventional interventions (e.g., acetaminophen, steroids, diphenhydramine, fluids, other pre-

medications medications or slowing of infusion rate) or a severe adverse event (anaphylaxis, anaphylactoid reactions, myocardial infarction, thromboembolism, or seizures) during or immediately after an infusion.

- The member is medically unstable (e.g. respiratory, cardiovascular, or renal conditions).
- The member has severe venous access issues that require the use of a special interventions only available in the outpatient hospital setting.
- The member has significant behavioral issues and/or physical or cognitive impairment that would impact the safety of the infusion therapy AND the patient does not have access to a caregiver.
- Alternative infusion sites (pharmacy, physician office, ambulatory care, etc.) are greater than 30 miles from the member's home.
- The member is less than 14 years of age.

For situations where administration of Leqembi does not meet the criteria for outpatient hospital infusion, coverage for Leqembi is provided when administered in alternative sites such as physician office, home infusion or ambulatory care.

Required Documentation

The following information is necessary to initiate the site of care prior authorization review (where applicable):

- Medical records supporting the member has experienced an adverse reaction that did not respond to conventional interventions or a severe adverse event during or immediately after an infusion
- Medical records supporting the member is medically unstable
- Medical records supporting the member has severe venous access issues that require specialized interventions only available in the outpatient hospital setting
- Medical records supporting the member has behavioral issues and/or physical or cognitive impairment and no access to a caregiver
- Records supporting alternative infusion sites are greater than 30 miles from the member's home
- Medical records supporting the member is new to therapy

Section 2: Clinical Criteria

Specialty Guideline Management Leqembi

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Leqembi	lecanemab-irmb

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications¹

Leqembi is indicated for the treatment of Alzheimer’s disease. Treatment with Leqembi should be initiated in patients with mild cognitive impairment or mild dementia stage of disease, the population in which treatment was initiated in clinical trials.

All other indications are considered experimental/investigational and not medically necessary.

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

Initial Requests

Genetic testing results documenting a mutation in amyloid precursor protein (APP), presenilin-1 (PSEN1), or presenilin-2 (PSEN2), if applicable.

Clinical documentation to support early onset Alzheimer’s Disease, if applicable.

Medical records (e.g., chart notes) documenting the following:

Diagnosis of Clinical Stage 3 or 4 Alzheimer’s Disease.

Baseline assessments for any of the following assessment tools:

Clinical Dementia Rating-Global Score (CDR-GS)

Mini-Mental Status Examination (MMSE)

Montreal Cognitive Assessment (MoCA)

Presence of amyloid pathology documented by either of the following:

- Baseline positron emission tomography (PET) scan
- Lumbar puncture results
- Recent (within one year) brain magnetic resonance imaging (MRI) prior to initiating

treatment.

Continuation requests (where applicable)

- Medical records (e.g., chart notes) documenting the most recent (less than 1 month prior to continuation request) assessment tool for any of the following:
 - Clinical Dementia Rating-Global Score (CDR-GS)
 - Mini-Mental Status Exam (MMSE)
 - Montreal Cognitive Assessment (MoCA)
- Brain magnetic resonance imaging (MRI) results within approximately one week prior to the 3rd, 5th dose, 7th dose, and 14th dose.

Prescriber Specialties

This medication must be prescribed by or in consultation with a geriatrician, neurologist, psychiatrist, or neuropsychiatrist.

Exclusions

- Coverage will not be provided for members with any of the following conditions:
 - suspected neurodegenerative etiology of cognitive impairment other than Alzheimer's disease (AD), including but not limited to frontotemporal lobar degeneration (FTLD) or Lewy body disease (i.e., meeting consensus criteria for possible or probable dementia with Lewy bodies that lack AD biomarkers of a positive amyloid PET or CSF profile).
 - > 4 cerebral microbleeds, cortical superficial siderosis, or a major vascular contribution to cognitive impairment confirmed via MRI.
 - Cerebral contusion, encephalomalacia, brain aneurysm or other vascular malformation, central nervous system infection, or brain tumor.
 - History of transient ischemic attacks (TIA), stroke, uncontrolled hypertension, or seizures within the past 12 months.
 - Bleeding disorder that is not under adequate control (including a platelet count <50,000 or international normalized ratio [INR] >1.5).
 - Immunologic disorder requiring therapy with immunoglobulins, monoclonal antibodies, immunosuppressants, or plasmapheresis.
- Leqembi will not be used in combination with any other amyloid beta-directed antibodies (e.g., aducanumab, donanemab).

Coverage Criteria

Alzheimer's Disease^{1,2,8-13,18-20}

Authorization of 7 months may be granted for treatment of Alzheimer's Disease (AD) when all of the following criteria are met:

- Member must meet one of the following criteria:
 - Member is 50 years of age or older
 - If less than 50 years of age, member has a genetic mutation in amyloid precursor protein (APP), presenilin-1 (PSEN1), or presenilin-2 (PSEN2), or other clinical documentation to support early onset AD.
- Member must have Clinical Stage 3 (cognitive impairment with early functional impact) or Clinical Stage 4 (dementia with mild functional impact) AD (Appendix A).
- Member must have objective evidence of cognitive impairment at baseline.
- Member must have one of the following scores at baseline on any of the following assessment tools:
 - Clinical Dementia Rating-Global Score (CDR-GS) of 0.5 or 1 (Appendix B).
 - Mini-Mental Status Examination (MMSE) score of 21 – 30 (Appendix C).
 - Montreal Cognitive Assessment (MoCA) score of greater than or equal to 16 (Appendix D).
- Member must meet one of the following criteria:
 - Have a positron emission tomography (PET) scan confirming the presence of amyloid pathology.
 - Have results from a lumbar puncture confirming at least one of the following detected in cerebrospinal fluid (CSF) as determined by the lab assay:
 - Low AB42/AB40 ratio
 - Elevated P-Tau/AB42 ratio
 - Elevated T-Tau/AB42 ratio
- Member must have a recent brain magnetic resonance imaging (MRI) within one year prior to initiating treatment to evaluate for pre-existing Amyloid Related Imaging Abnormalities (ARIA).
- Member will not use the requested medication in combination with anticoagulants including warfarin, heparin, and direct oral anticoagulants (e.g., dabigatran, rivaroxaban, edoxavan, apiximab, betrixaban).
- If there is concurrent use of antiplatelet agents (e.g., aspirin up to 325 mg/day, clopidogrel, prasugrel, ticagrelor), member will use antiplatelet agent as monotherapy at a standard therapeutic dose (i.e., not using as dual agent anti-platelet therapy).
- Member has undergone genotype testing to determine apolipoprotein E ε4 (ApoE ε4) status prior to initiation of the requested medication to inform member of the risk of developing ARIA.:
- Member and/or provider must currently be participating in a provider-enrolled patient registry that collects information on treatments for Alzheimer's disease (e.g., Alzheimer's Network for Treatment and Diagnostics (ALZ-NET)).

Continuation of Therapy

Authorization of 12 months (first reauthorization after the initial 7-month approval period) may be granted for members requesting continuation of therapy when all of the following criteria are met:

- Member has met all requirements in the coverage criteria at the time of initial approval.
- Member has been evaluated for evidence of amyloid-related imaging abnormalities (ARIA) on MRI within approximately one week prior to the 3rd, 5th dose, 7th dose, and 14th dose (Appendix E).
 - For members with radiographic evidence of ARIA-E:
 - Dosing may continue based on clinical judgement, if applicable, for members that meet the following criteria:
 - Member has mild ARIA-E on MRI and is asymptomatic or has mild clinical symptoms
 - Dosing should be suspended until MRI demonstrates radiographic resolution and symptoms resolve for members that meet any of the following criteria:
 - Member has mild ARIA-E on MRI and has moderate or severe clinical symptoms
 - Member has moderate ARIA-E on MRI and is asymptomatic or has mild, moderate, or severe clinical symptoms
 - Member has severe ARIA-E on MRI and is asymptomatic or has mild, moderate, or severe clinical symptoms
 - For members with radiographic evidence of ARIA-H:
 - Dosing may continue for members that meet the following criteria:
 - Member has mild ARIA-H on MRI and is asymptomatic
 - Dosing should be suspended until MRI demonstrates radiographic stabilization and symptoms resolve for members that meet any of the following criteria:
 - Member has mild ARIA-H on MRI and is symptomatic
 - Member has moderate ARIA-H on MRI and is asymptomatic or symptomatic
 - Member has severe ARIA-H on MRI and is asymptomatic or symptomatic
- Member and/or provider continues to participate in a provider-enrolled patient registry that collects information on treatments for Alzheimer's disease (e.g., Alzheimer's Network for Treatment and Diagnostics (ALZ-NET)).

Authorization of 12 months (reauthorizations beyond initial 19 months of therapy) may be granted for members requesting continuation of therapy when all of the following criteria are met:

- Member has met all requirements in the coverage criteria at the time of initial approval.
- Member has a positive clinical response as evidenced by stabilization or slowing of disease progression as documented by any of the following (Note: repeat assessment tool(s) must be the same tool that was submitted upon initial request):
 - CDR-Global Score (i.e., score of 0.5 or 1)

- MMSE (i.e., decline of 3 points or less per year)
- MoCA (i.e., score of greater than or equal to 16)

Note: Continuation requests for members with assessment scores outside of the provided ranges (i.e. mild dementia) or who have progressed greater than the provided rate of decline may be reviewed on a case-by-case basis.

- Member and/or provider continues to participate in a provider-enrolled patient registry that collects information on treatments for Alzheimer’s disease (e.g., Alzheimer’s Network for Treatment and Diagnostics (ALZ-NET)).

Appendix

Appendix A: Diagnostic criteria for mild cognitive impairment (MCI) and dementia with mild functional impact^{14,19-20}

- Mild cognitive impairment (MCI)/Clinical Stage 3:
 - Cognitive concerns by the patient, knowledgeable informant, or the physician
 - Objective impairment in one or more cognitive domains including memory, executive function, attention, language, or visuospatial skills
 - Generally preserved activities of daily living (ADL)
 - No dementia
- Dementia with mild functional impairment/Clinical Stage 4:
 - Cognitive concerns by the patient, knowledgeable informant, or the physician
 - Performance in the impaired/abnormal range on objective cognitive tests
 - Evidence of decline from baseline, documented by the individual’s report or by observer (e.g., study partner) report or by change on longitudinal cognitive testing or neurobehavioral assessments
 - Progressive cognitive and mild functional impairment on instrumental ADL with independence in basic ADL

Appendix B: Clinical Dementia Rating (CDR) Scale³⁻⁵

The CDR is obtained through semi-structured interviews of patients and informants with cognitive functioning rated on a 5-point scale in the following domains: memory, orientation, judgment and problem solving, community affairs, home and hobbies, and personal care. The score relates to the member’s level of dementia:

- 0 = Normal
- 0.5 = Very Mild Dementia
- 1 = Mild Dementia
- 2 = Moderate Dementia
- 3 = Severe Dementia

Appendix C: Mini-Mental Status Exam (MMSE)⁶

The MMSE is scored on a 30-point scale, with items that assess orientation (temporal and spatial; 10 points), memory (registration and recall; 6 points), attention/concentration (5 points), language (verbal and written, 8 points), and visuospatial function (1 point). The score relates to the member’s level of dementia:

- 25 – 30 suggests normal cognition
- 20 – 24 suggests mild dementia
- 13 – 20 suggests moderate dementia
- Less than 12 suggests severe dementia

Appendix D: Montreal Cognitive Assessment (MoCA)^{7,8}

Per MoCA assessment, average scores for the following ranges are:

- Mild Cognitive Impairment: 19 – 25
- Mild Dementia: 11 – 21
- Normal: 26 and above

Appendix E: ARIA MRI Classification Criteria¹

ARIA Type	Radiographic Severity Mild	Radiographic Severity Moderate	Radiographic Severity Severe
ARIA-E	FLAIR hyperintensity confined to sulcus and or cortex/subcortical white matter in one location < 5 cm	FLAIR hyperintensity 5 to 10 cm, or more than 1 site of involvement, each measuring < 10 cm	FLAIR hyperintensity measuring > 10 cm with associated gyral swelling and sulcal effacement. One or more separate/independent sites of involvement may be noted.
ARIA-H microhemorrhage	≤ 4 new incident microhemorrhages	5 to 9 new incident microhemorrhages	10 or more new incident microhemorrhages
ARIA-H superficial siderosis	1 focal area of superficial siderosis	2 focal areas of superficial siderosis	> 2 areas of superficial siderosis

REFERENCES

SECTION 1

1. Leqembi [package insert]. Nutley, NJ: Eisai, Inc.; January 2025.

SECTION 2

2. Leqembi [package insert]. Nutley, NJ: Eisai Inc.; January 2025.
3. Fagan AM, Mintun MA, Mach RH, et al. Inverse relation between in vivo amyloid imaging load and cerebrospinal fluid Abeta42 in humans. *Ann Neurol*. 2006;59(3):512-519.
4. O'Bryant SE, Waring SC, Cullum CM, et al. Staging dementia using Clinical Dementia Rating Scale Sum of Boxes scores: a Texas Alzheimer's research consortium study. *Arch Neurol*. 2008;65(8):1091-1095. Doi:10.1001/archneur.65.8.1091.
5. CDR Dementia Staging Instrument. Knight Alzheimer Disease Research Center. <https://knightadrc.wustl.edu/cdr/cdr.htm>. Accessed: May 1, 2024.
6. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology*. 1993 Nov;43(11):2412-4.
7. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975 Nov;12(3):189-98.
8. MoCA Cognitive Assessment. <https://www.mocatest.org/>. Accessed: May 4, 2023.
9. Nasreddine ZS, Phillips NA, Bédirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment [published correction appears in *J Am Geriatr Soc*. 2019 Sep;67(9):1991]. *J Am Geriatr Soc*. 2005;53(4):695-699.
10. Schindler SE, Gray JD, Gordon BA, et al. Cerebrospinal fluid biomarkers measured by Elecsys assays compared to amyloid imaging. *Alzheimers Dement*. 2018;14(11):1460-1469.
11. Cummings J, Aisen P, Apostolova LG, Atri A, Salloway S, Weiner M. Aducanumab: Appropriate Use Recommendations. *J Prev Alzheimers Dis*. 2021;8(4):398-410.
12. Patrick RE, Hobbs K, Mathias L, Harper DG, Forester BP. The Limitations of Using Cognitive Cutoff Scores for Enrollment in Alzheimer Trials. *Am J Geriatr Psychiatry*. 2019;27(10):1153-1158.
13. National Coverage Determination (NCD) for Monoclonal Antibodies Directed Against Amyloid for the Treatment of Alzheimer's Disease (AD) (200.3 – Version 1). <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=375&ncdver=1> Accessed May 1, 2025.
14. Trzepacz PT, Hochstetler H, Wang S, Walker B, Saykin AJ; Alzheimer's Disease Neuroimaging Initiative. Relationship between the Montreal Cognitive Assessment and Mini-mental State Examination for assessment of mild cognitive impairment in older adults. *BMC Geriatr*. 2015 Sep 7;15:107.
15. Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011;7(3):270-279.
16. Clark CM, Sheppard L, Fillenbaum GG, Galasko D, et al. Variability of annual Mini-Mental State Examination score in patients with probable Alzheimer disease: a clinical perspective of data from the Consortium to Establish a Registry for Alzheimer's Disease. *Arch Neurol*. 1999 Jul;56(7):857-62.
17. Han L, Cole M, Bellavance F, McCusker J, Primeau F. Tracking cognitive decline in Alzheimer's disease using the mini-mental state examination: a meta-analysis. *Int Psychogeriatr*. 2000 Jun;12(2):231-47.
18. Morris JC, Edland S, Clark C, Galasko D, et al. The consortium to establish a registry for Alzheimer's disease (CERAD). Part IV. Rates of cognitive change in longitudinal assessment of probable Alzheimer's disease. *Neurology*. 1993 Dec;43(12):2457-65.

19. Elecsys Phospho-Tau (181P) CSF 2022-12.
20. Rabinovici GD, Selkoe DJ, Schindler SE, et al. Donanemab: Appropriate use recommendations. *J Prev Alzheimers Dis*. Published online March 27, 2025.
21. Jack CR Jr, Andrews JS, Beach TG, et al. Revised criteria for diagnosis and staging of Alzheimer's disease: Alzheimer's Association Workgroup. *Alzheimers Dement*. 2024;20(8):5143-5169.