

POLICY Document for octreotide products

The overall objective of this policy is to support the appropriate and cost-effective use of the medication, specific to use of preferred medication options, and overall, clinically appropriate use. This document provides specific information to both sections of the overall policy.

Section 1: Clinical Criteria

- Policy information specific to the clinical appropriateness for the medication

Section 2: Oncology Clinical Policy

- Policy information specific to regimen review per NCCN Guidelines.

Section 1: Clinical Criteria

Specialty Guideline Management octreotide products

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Sandostatin	octreotide acetate injection
Bynfezia Pen	octreotide acetate injection
Mycapssa	octreotide delayed-release capsule
Sandostatin LAR Depot	octreotide acetate for injectable suspension
octreotide acetate injection	octreotide acetate injection
octreotide acetate for injectable suspension	octreotide acetate for injectable suspension

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

- FDA-approved Indications

- Sandostatin/Bynfezia Pen/ octreotide acetate injection:
 - Indicated to reduce blood levels of growth hormone (GH) and insulin growth factor-1 (IGF-1; somatomedin C) in acromegaly patients who have had inadequate response to or cannot be treated with surgical resection, pituitary irradiation, and bromocriptine mesylate at maximally tolerated doses.
 - Indicated for the symptomatic treatment of patients with metastatic carcinoid tumors where it suppresses or inhibits the severe diarrhea and flushing episodes associated with the disease.
 - Indicated for the treatment of the profuse watery diarrhea associated with vasoactive intestinal peptide (VIP)-secreting tumors.
- Sandostatin LAR/octreotide acetate for injectable suspension: Sandostatin LAR Depot and octreotide acetate for injectable suspension are indicated in patients in whom initial treatment with Sandostatin Injection/octreotide acetate injection have been shown to be effective and tolerated.
 - Long-term maintenance therapy in acromegalic patients who have had an inadequate response to surgery and/or radiotherapy, or for whom surgery and/or radiotherapy is not an option.
 - Long-term treatment of the severe diarrhea and flushing episodes associated with metastatic carcinoid tumors.
 - Long-term treatment of the profuse watery diarrhea associated with vasoactive intestinal peptide (VIP)-secreting tumors.
- Mycapssa is indicated for long-term maintenance treatment in acromegaly patients who have responded to and tolerated treatment with octreotide or lanreotide.

• **Compendial Uses (applies to injectable products)**

- Neuroendocrine tumors (NETs):
 - Tumors of the gastrointestinal (GI) tract, lung, and thymus (carcinoid tumors)
 - Tumors of the pancreas (islet cell tumors)
 - Gastroenteropancreatic neuroendocrine tumors (GEP-NETs)
- Pheochromocytoma and paraganglioma
- Thymomas and thymic carcinomas
- Congenital hyperinsulinism (CHI)/persistent hyperinsulinemic hypoglycemia of infancy (PHHI)(octreotide and Sandostatin only)
- Acquired immune deficiency syndrome (AIDS)-associated diarrhea
- Inoperable bowel obstruction
- Cancer-related diarrhea
- Enterocutaneous fistula
- Gastroesophageal varices
- Pancreatic fistulas
- Pituitary adenoma
- Short bowel syndrome
- Zollinger-Ellison syndrome

All other indications are considered experimental/investigational and not medically necessary.

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

- For acromegaly:
 - For initial approval: Laboratory report indicating high pretreatment insulin-like growth factor-1 (IGF-1) level and chart notes indicating an inadequate or partial response to surgery or radiotherapy or a clinical reason for not having surgery or radiotherapy.
 - For continuation: Laboratory report indicating normal current IGF-1 levels or chart notes indicating that the member's IGF-1 level has decreased or normalized since initiation of therapy.
- Cancer-related diarrhea: Chart notes indicating grade 3 or 4 diarrhea.

Coverage Criteria

- **Acromegaly**

Authorization of 12 months may be granted for the treatment of acromegaly when all of the following criteria are met:

- Member has a high pretreatment IGF-1 level for age and/or gender based on the laboratory reference range.
- Member had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason why the member has not had surgery or radiotherapy.
- For Mycapssa requests, member has previously responded to and tolerated treatment with octreotide or lanreotide.

- **Neuroendocrine Tumors (NETs) (injectable products only)**

- Authorization of 12 months may be granted for treatment of NETs of the gastrointestinal (GI) tract, lung, and thymus (carcinoid tumors).
- Authorization of 12 months may be granted for treatment of NETs of the pancreas (islet cell tumors), including gastrinomas, glucagonomas, and insulinomas.
- Authorization of 12 months may be granted for treatment of gastroenteropancreatic neuroendocrine tumors (GEP-NETs).

- **Carcinoid Syndrome (injectable products only)**

Authorization of 12 months may be granted for treatment of carcinoid syndrome.

- **Vasoactive Intestinal Peptide Tumors (VIPomas) (injectable products only)**

Authorization of 12 months may be granted for management of symptoms related to hormone hypersecretion of VIPomas.

- **Pheochromocytoma and Paraganglioma (injectable products only)**

Authorization of 12 months may be granted for treatment of pheochromocytoma and paraganglioma.

- **Thymomas and Thymic Carcinomas (injectable products only)**

Authorization of 12 months may be granted for treatment of thymomas and thymic carcinomas.

- **Congenital Hyperinsulinism (CHI)/Persistent Hyperinsulinemic Hypoglycemia of Infancy (octreotide and Sandostatin only)**

Authorization of 6 months may be granted for treatment of CHI and persistent hyperinsulinemic hypoglycemia in an infant.

- **AIDS-Associated Diarrhea (injectable products only)**

Authorization of 12 months may be granted for treatment of AIDS-associated severe secretory diarrhea when anti-microbial (e.g., ciprofloxacin or metronidazole) or anti-motility agents (e.g., loperamide or diphenoxylate and atropine) have become ineffective.

- **Inoperable Bowel Obstruction in Cancer (injectable products only)**

Authorization of 12 months may be granted for management of GI symptoms (e.g., nausea, pain, vomiting) of inoperable bowel obstruction in members with cancer.

- **Cancer-Related Diarrhea (injectable products only)**

Authorization of 12 months may be granted for treatment of cancer-related diarrhea when the member has grade 3 or greater diarrhea according to National Cancer Institute (NCI) Common Terminology Criteria for Adverse Events (CTCAE).

- **Enterocutaneous Fistula (injectable products only)**

Authorization of 12 months may be granted for management of volume depletion from enterocutaneous fistula.

- **Gastroesophageal Varices (injectable products only)**

Authorization of 6 months may be granted for treatment of acute bleeding of gastroesophageal varices associated with cirrhosis.

- **Pancreatic Fistulas (injectable products only)**

Authorization of 6 months may be granted for prevention and treatment of pancreatic fistulas following pancreatic surgery.

- **Pituitary Adenoma (injectable products only)**

Authorization of 12 months may be granted for treatment of pituitary adenoma.

- **Short Bowel Syndrome (injectable products only)**

Authorization of 12 months may be granted for treatment of short bowel syndrome when the daily intravenous fluid requirement is greater than 3 liters.

- **Zollinger-Ellison Syndrome (injectable products only)**

Authorization of 12 months may be granted for treatment of Zollinger-Ellison syndrome.

Continuation of Therapy

- **Acromegaly**

Authorization of 12 months may be granted for continuation of therapy for acromegaly when the member's IGF-1 level has decreased or normalized since initiation of therapy.

- **NETs, Carcinoid Syndrome, Vipomas, Pheochromocytoma/Paraganglioma, Thymomas/Thymic Carcinomas, AIDS-Associated Diarrhea, Bowel Obstruction, Cancer-Related Diarrhea, And Zollinger-Ellison Syndrome (injectable products only)**

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization when the member is experiencing clinical benefit as evidenced by improvement or stabilization in clinical signs and symptoms since initiation of therapy.

- **All Other Indications**

All members (including new members) requesting authorization for continuation of therapy must meet all requirements in the coverage criteria.

Section 2: Oncology Clinical Policy

PURPOSE

The purpose of this policy is to define the Novologix NCCN® Regimen Prior Authorization Program.

SCOPE

This policy applies to clients who have implemented the Novologix NCCN® Program as a part of their medical and/or pharmacy prior authorization solution.

PROGRAM DESCRIPTION

The National Comprehensive Care Network® (NCCN®) is an alliance of leading cancer centers devoted to patient care, research and education dedicated to improving the quality, effectiveness, and efficiency of cancer care so patients can live better lives.¹ It is comprised of oncology experts who convene regularly to establish the best treatments for patients. NCCN develops various resources for use by stakeholders in the health care delivery system. These resources include, but are not limited to, the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®), the NCCN Drugs & Biologics Compendium (NCCN Compendium®) and the NCCN Chemotherapy Order Templates (NCCN Templates®).

NCCN Templates® are based on NCCN Guidelines® and NCCN Compendium®. The NCCN Compendium lists the appropriate drugs and biologics as treatment options for specific cancers using U.S. Food and Drug Administration (FDA)-approved disease indications and specific NCCN panel recommendations. Each recommendation is supported by a level of evidence category.

NCCN Categories of Evidence and Consensus²

- Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
- Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
- Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
- Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

POLICY

Policy for Regimen Prior Authorization

octreotide products SGM 1734-A P2024a_R.docx
Novologix LLC_NCCN Oncology Clinical Policy_9.2024

© 2024 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

A regimen prior authorization allows submission of a single prior authorization request for all oncology drugs or biologics within an NCCN template that require prior authorization.

PROCEDURE

This policy provides coverage of a regimen review when all of the following criteria are met:

1. Regimen prior authorization reviews, based on NCCN templates, are initiated through the provider portal.
 - If the prior authorization request is submitted via phone or fax, each drug or biologic will need to be submitted and reviewed as a separate prior authorization request for review with drug-specific criteria.
2. The prior authorization review is requested for an oncology drug or biologic.
3. The member is eligible for regimen review.
4. The indication is for a cancer that is eligible for regimen review. Currently, the cancer types in scope for regimen review include the following:
 - o Ampullary Adenocarcinoma
 - o Anal Carcinoma
 - o B-Cell Lymphomas
 - o Basal Cell Skin Cancer
 - o Biliary Tract Cancers
 - o Bone Cancer
 - o Breast Cancer
 - o Bladder Cancer
 - o Central Nervous System Cancers
 - o Cervical Cancer
 - o Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma
 - o Chronic Myeloid leukemia
 - o Colon Cancer
 - o Dermatofibrosarcoma Protuberans
 - o Esophageal Cancer
 - o Gastric Cancer
 - o Gastrointestinal Stromal Tumors
 - o Gestational Trophoblastic Neoplasms
 - o Hairy Cell Leukemia
 - o Head and Neck Cancers
 - o Histiocytic Neoplasms
 - o Hodgkin Lymphoma
 - o Hepatocellular Carcinoma
 - o Kaposi Sarcoma
 - o Kidney Cancer

- o Melanoma: Cutaneous
- o Melanoma: Uveal
- o Merkel Cell Carcinoma
- o Mesothelioma: Peritoneal
- o Mesothelioma: Pleural
- o Multiple Myeloma
- o Myelodysplastic Syndromes
- o Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions
- o Myeloproliferative Neoplasms
- o Neuroendocrine and Adrenal Tumors
- o Non-Small Cell Lung Cancer
- o Occult Primary
- o Ovarian Cancer
- o Pancreatic Cancer
- o Penile Cancer
- o Primary Cutaneous Lymphomas
- o Prostate Cancer
- o Rectal Cancer
- o Small Bowel Adenocarcinoma
- o Small Cell Lung Cancer
- o Soft Tissue Sarcoma
- o Squamous Cell Skin Cancer
- o Systemic Mastocytosis
- o Systemic Light Chain Amyloidosis
- o T-Cell Lymphomas
- o Testicular Cancer
- o Thymomas and Thymic Carcinomas
- o Thyroid Carcinoma
- o Uterine Neoplasms
- o Vaginal Cancer
- o Vulvar Cancer
- o Waldenström Macroglobulinemia / Lymphoplasmacytic Lymphoma
- o Wilms Tumor (Nephroblastoma)

In addition, the following criteria must be met for approval:

1. The requested regimen for the drug(s) or biologic(s) and indication is consistent with an NCCN recommendation with a level of evidence category of 1 or 2A.
2. The NCCN template must be accepted by the provider without modification.

Further review may be indicated when the above criteria are not met.

Authorizations may be granted for 12 months or as medically required, based on the member's condition and provider's assessment.

Supportive Care: Myeloid Growth Factor Therapy

Granulocyte colony stimulating factors are recommended for primary prophylaxis based on the febrile neutropenia risk of the chemotherapy regimen. Febrile neutropenia risk levels vary by NCCN Chemotherapy Order template and are listed at the top of the template. Regimens associated with a high or intermediate risk of febrile neutropenia may include a granulocyte colony stimulating factor as part of the prior authorization.

Continuation of Therapy

To submit a request for continuation of therapy, a new regimen prior authorization review must be requested. Upon template selection, the template must be modified to include the appropriate therapies being used for maintenance treatment. The regimen request will be submitted for further review.

Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and evidence-based practice guidelines.

REFERENCES:

SECTION 1

1. Octreotide acetate [package insert]. Morgantown, WV: Mylan Institutional LLC; November 2022.
2. Sandostatin [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; November 2023.
3. Sandostatin LAR Depot [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; July 2023.
4. The NCCN Drugs & Biologics Compendium® © 2023 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed November 14, 2023.
5. Clinical Consult. CVS Caremark Clinical Programs Review: Focus on Pediatric Endocrinology. September 2007.
6. Katznelson L, Laws ER, Melmed S, et al. Acromegaly: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2014;99:3933-3951.
7. American Association of Clinical Endocrinologists Acromegaly Guidelines Task Force. Medical guidelines for clinical practice for the diagnosis and treatment of acromegaly – 2011 update. *Endocr Pract*. 2011;17(suppl 4):1-44.
8. The NCCN Clinical Practice Guidelines in Oncology® Neuroendocrine and Adrenal Tumors (Version 1.2023). © 2023 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed November 14, 2023.

9. Rinke A, Muller H, Schade-Brittinger C, et al. Placebo-controlled, double-blind, prospective, randomized study on the effect of octreotide LAR in the control of tumor growth in patients with metastatic neuroendocrine midgut tumors: a report from the PROMID study group. *J Clin Oncol*. 2009;27:4656-4663.
10. Thapar K, Kovacs K, Stefaneanu L, et al: Antiproliferative effect of the somatostatin analogue octreotide on growth hormone-producing pituitary tumors: Results of a multicenter randomized trial. *Mayo Clin Proc* 1997; 72:893-900.
11. The NCCN Clinical Practice Guidelines in Oncology® Thymomas and Thymic Carcinomas. (Version 1.2023). © 2023 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed November 14, 2023.
12. The NCCN Clinical Practice Guidelines in Oncology® Palliative Care (Version 2.2023). © 2023 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed November 14, 2023.
13. Harris AG, O'Dorisio TM, Woltering EA, et al. Consensus statement: Octreotide dose titration in secretory diarrhea. Diarrhea Management Consensus Development Panel. *Dig Dis Sci*. 1995;40(7):1464-1473.
14. Fried M. Octreotide in the treatment of refractory diarrhea. *Digestion*. 1999;60:42-46.
15. Edmunds MC, Chen JD, Soykan I, et al. Effect of octreotide on gastric and small bowel motility in patients with gastroparesis. *Aliment Pharmacol Ther*. 1998;12(2):167-174.
16. Ripamonti C, Mercadante S, Groff L, et al. Role of octreotide, scopolamine butylbromide, and hydration in symptom control of patients with inoperable bowel obstruction and nasogastric tubes: A prospective randomized trial. *J Pain Symptom Manage*. 2000;19(1):23-34.
17. Mercadante S, Ripamonti C, Casuccio A, et al. Comparison of octreotide and hyoscine butylbromide in controlling gastrointestinal symptoms due to malignant inoperable bowel obstruction. *Support Care Cancer*. 2000;8(3):188-191.
18. Dorta G. Role of octreotide and somatostatin in the treatment of intestinal fistulae. *Digestion*. 1999;60 Suppl 2:53-56.
19. Jamil M, Ahmed U, Sobia H. Role of somatostatin analogues in the management of enterocutaneous fistulae. *J Coll Physicians Surg Pak*. 2004;14(4):237-240.
20. Rahbour G, Siddiqui MR, Ullah MR, et al. A meta-analysis of outcomes following use of somatostatin and its analogues for the management of enterocutaneous fistulas. *Ann Surg*. 2012;256(6):946-954.
21. Freitas DS, Sofia C, Pontes JM, et al. Octreotide in acute bleeding esophageal varices: A prospective randomized study. *Hepatogastroenterology*. 2000;47(35):1310-1314.
22. Imperiale TF, Teran JC, McCullough AJ. A meta-analysis of somatostatin versus vasopressin in the management of acute esophageal variceal hemorrhage. *Gastroenterology*. 1995;109(4):1289-1294.
23. Gøtzsche PC, Hróbjartsson A. Somatostatin analogues for acute bleeding oesophageal varices. *Cochrane Database Syst Rev*. 2008;(3):CD000193.
24. Erstad BL. Octreotide for acute variceal bleeding. *Ann Pharmacother*. 2001;35(5):618-626.
25. Corley DA, Cello JP, Adkisson W, et al. Octreotide for acute esophageal variceal bleeding: A meta-analysis. *Gastroenterology*. 2001;120(4):946-954.
26. Gross M, Schiemann U, Muhlhofer A, Zoller WG. Meta-analysis: Efficacy of therapeutic regimens in ongoing variceal bleeding. *Endoscopy*. 2001;33(9):737-746.

27. Li-Ling J, Irving M. Somatostatin and octreotide in the prevention of postoperative pancreatic complications and the treatment of enterocutaneous pancreatic fistulas: A systematic review of randomized controlled trials. *Br J Surg*. 2001;88(2):190-199.
28. Machado NO. Pancreatic fistula after pancreatectomy: Definitions, risk factors, preventive measures, and management – Review. *Int J Surg Oncol*. 2012;2012:602478.
29. Adachi T, Kuroki T, Kitasato A, et al. Safety and efficacy of early drain removal and triple-drug therapy to prevent pancreatic fistula after distal pancreatectomy. *Pancreatology* 2015;15:411-416.
30. Gurusamy KS, Koti R, Fusai G, Davidson BR. Somatostatin analogues for pancreatic surgery. *Cochrane Database Syst Rev*. 2013;4:CD008370.
31. Alberta Provincial CNS Tumour Team. Pituitary adenomas. Clinical Practice Guideline No. CNS-006 Edmonton, AB: Alberta Health Services, Cancer Care; August 2012.
32. Peeters M, Van den Brande J, Francque S. Diarrhea and the rationale to use Sandostatin. *Acta Gastroenterol Belg*. 2010;73(1):25-36.
33. Mystakidou K, Tsilika E, Kalaidopoulou O, et al. Comparison of octreotide administration vs conservative treatment in the management of inoperable bowel obstruction in patients with far advanced cancer: A randomized, double-blind, controlled clinical trial. *Anticancer Res*. 2002;22(2B):1187-1192.
34. American Gastroenterological Association. American Gastroenterological Association medical position statement: Short bowel syndrome and intestinal transplantation. *Gastroenterology*. 2003;124(4):1105-1110.
35. Loehrer PJ Sr, Wang W, Johnson DH, et al. Octreotide alone or with prednisone in patients with advanced thymoma and thymic carcinoma: An Eastern Cooperative Oncology Group Phase II Trial. *J Clin Oncol*. 2004;22(2):293-299.
36. Leandros E, Antonakis PT, Albanopoulos K, et al. Somatostatin versus octreotide in the treatment of patients with gastrointestinal and pancreatic fistulas. *Can J Gastroenterol*. 2004;18(5):303-306.
37. Allen PJ, Gonen M, Brennan MF, et al. Pasireotide for postoperative pancreatic fistula. *N Engl J Med*. 2014;370(21):2014-2022.
38. Bynfezia Pen [package insert]. Cranbury, NJ: Sun Pharmaceutical Industries Inc.; April 2020.
39. Mycapssa [package insert]. Dublin, Ireland: Amryt Pharmaceuticals, Inc.; March 2022.
40. Lexicomp Online®, Lexi-Drugs. Waltham, MA: UpToDate, Inc.; Updated November 1, 2023. <http://online.lexi.com> [available with subscription]. Accessed November 10, 2023.
41. Octreotide acetate for injectable suspension. Parsippany, NJ: Teva Pharmaceuticals; January 2024.

SECTION 2

1. National Comprehensive Cancer Network. About NCCN website. <https://www.nccn.org/home/about>, accessed September 9, 2024.
2. National Comprehensive Cancer Network. NCCN Categories of Evidence and Consensus website, <https://www.nccn.org/guidelines/guidelines-process/development-and-update-of-guidelines>, accessed September 9, 2024.

3. National Comprehensive Cancer Network. NCCN Guidelines website. https://www.nccn.org/guidelines/category_1, accessed September 9, 2024. (Note: An account may be required.)
4. National Comprehensive Cancer Network. NCCN Drugs and Biologics Compendium website <https://www.nccn.org/compendia-templates/compendia/drugs-and-biologics-compendia>, accessed September 9, 2024. (Note: A subscription may be required.)
5. National Comprehensive Cancer Network. NCCN Chemotherapy Order Templates (NCCN Templates) website. <https://www.nccn.org/compendia-templates/nccn-templates-main/browse-by-cancer-type>, accessed September 9, 2024. (Note: A subscription may be required.)