



Pre - PA Allowance

None

Prior-Approval Requirements

Age 18 years of age or older

Diagnosis

Patient must have the following:

1. Metastatic non-small cell lung cancer (NSCLC)
 - a. Anaplastic lymphoma kinase (ALK)-positive
 - b. Female patients of reproductive potential **only**: patient will be advised to use effective non-hormonal contraception during treatment with Alunbrig and for 4 months after the final dose
 - c. Males with female partners of reproductive potential **only**: patient will be advised to use effective contraception during treatment with Alunbrig and for 3 months after the final dose

Prior - Approval Limits

Quantity

Strength	Quantity
30 mg	540 tablets per 90 days OR
90 mg	180 tablets per 90 days OR
One Month Initiation Pack (7 tabs of 90mg & 23 tabs of 180mg) + 180 mg	1 Initiation Pack + 90 tablets per 90 days OR
180 mg	90 tablets per 90 days

Maximum daily limit of any combination: 180 mg

*** Quantity limits listed above must be used to achieve dose optimization**

****Utilizing the highest strengths available to achieve the dosage is recommended to minimize dosing errors and improve compliance**

Duration 12 months

Prior – Approval *Renewal* Requirements

Age 18 years of age or older

Diagnosis



**ALUNBRIG
(brigatinib)**

Patient must have the following:

1. Metastatic non-small cell lung cancer (NSCLC)
 - a. **NO** disease progression or unacceptable toxicity
 - b. Female patients of reproductive potential **only**: patient will be advised to use effective non-hormonal contraception during treatment with Alunbrig and for 4 months after the final dose
 - c. Males with female partners of reproductive potential **only**: patient will be advised to use effective contraception during treatment with Alunbrig and for 3 months after the final dose

Prior - Approval *Renewal* Limits

Quantity

Strength	Quantity
30 mg	360 tablets per 90 days OR
90 mg	180 tablets per 90 days OR
180 mg	90 tablets per 90 days

Maximum daily limit of any combination: 180 mg

*** Quantity limits listed above must be used to achieve dose optimization**

****Utilizing the highest strengths available to achieve the dosage is recommended to minimize dosing errors and improve compliance**

Duration 12 months