

**BRUKINSA  
(zanubrutinib)**

**Pre - PA Allowance**

None

---

**Prior-Approval Requirements**

**Age** 18 years of age or older

**Diagnoses**

Patient must have **ONE** of the following:

1. Mantle cell lymphoma (MCL)
  - a. Patient has received at least one prior therapy
2. Waldenström's macroglobulinemia (WM)
3. Relapsed or refractory marginal zone lymphoma (MZL)
  - a. Patient has received at least one anti-CD20-based regimen
4. Chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL)
5. Relapsed or refractory follicular lymphoma (FL)
  - a. Patient has received two or more lines of systemic therapy
  - b. Used in combination with Gazyva (obinutuzumab)

**AND ALL** of the following:

- a. Prescriber agrees to monitor for bleeding and malignancies
- b. Prescriber agrees to monitor CBC for cytopenias
- c. Prescriber agrees to monitor for cardiac arrhythmias
- d. Females of reproductive potential **only**: patient will be advised not to become pregnant during treatment with Brukinsa and for at least 1 week after the last dose
- e. Males with female partners of reproductive potential **only**: patient will be advised not to father a child during treatment with Brukinsa and for at least 1 week after the last dose

**Prior - Approval Limits**

**Quantity** 360 capsules per 90 days

**Duration** 12 months

---

**Prior – Approval *Renewal* Requirements**

**Age** 18 years of age or older



**BlueCross  
BlueShield**

Federal Employee Program.

## **BRUKINSA (zanubrutinib)**

### **Diagnoses**

Patient must have **ONE** of the following:

1. Mantle cell lymphoma (MCL)
2. Waldenström's macroglobulinemia (WM)
3. Relapsed or refractory marginal zone lymphoma (MZL)
4. Chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL)
5. Relapsed or refractory follicular lymphoma (FL)
  - a. Used in combination with Gazyva (obinutuzumab)

**AND ALL** of the following:

- a. **NO** disease progression or unacceptable toxicity
- b. Prescriber agrees to monitor for bleeding and malignancies
- c. Prescriber agrees to monitor CBC for cytopenias
- d. Prescriber agrees to monitor for cardiac arrhythmias
- e. Females of reproductive potential **only**: patient will be advised not to become pregnant during treatment with Brukinsa and for at least 1 week after the last dose
- f. Males with female partners of reproductive potential **only**: patient will be advised not to father a child during treatment with Brukinsa and for at least 1 week after the last dose

### **Prior - Approval *Renewal* Limits**

Same as above