

## Pre - PA Allowance

None

---

## Prior-Approval Requirements

**Age** 18 years of age or older

### Diagnosis

Patient must have the following:

Newly-diagnosed acute myeloid leukemia (AML)

**AND ALL** of the following:

1. Used in combination with low-dose cytarabine
2. Patient is 75 years of age or older **OR** patient has comorbidities that preclude the use of intensive induction chemotherapy
3. Prescriber agrees to monitor electrocardiograms (ECGs) for QTc prolongation
4. Prescriber agrees to advise females of reproductive potential to use effective contraception during treatment and for at least 30 days after the last dose
5. Prescriber agrees to advise males with female partners of reproductive potential to use condoms during treatment and for at least 30 days after the last dose

## Prior - Approval Limits

### Quantity

Strength	Quantity Limit
25 mg tablets	180 tablets per 90 days <b>OR</b>
100 mg tablets	90 tablets per 90 days

**Duration** 12 months

---

## Prior – Approval *Renewal* Requirements

**Age** 18 years of age or older

### Diagnosis

Patient must have the following:

Acute myeloid leukemia (AML)

**AND ALL** of the following:

1. Used in combination with low-dose cytarabine
2. **NO** disease progression or unacceptable toxicity
3. Prescriber agrees to monitor electrocardiograms (ECGs) for QTc prolongation
4. Prescriber agrees to advise females of reproductive potential to use effective contraception during treatment and for at least 30 days after the last dose
5. Prescriber agrees to advise males with female partners of reproductive potential to use condoms during treatment and for at least 30 days after the last dose

## **Prior - Approval *Renewal* Limits**

Same as above