

## Pre - PA Allowance

None

---

## Prior-Approval Requirements

**Age** 18 years of age or older

### Diagnoses

Patient must have **ONE** of the following:

1. Squamous cell carcinoma of the head and neck
  - a. Stage III
    - i. If non-nasopharyngeal site- concurrent radiation therapy
  - b. Stage IV
    - i. If non-nasopharyngeal site- concurrent radiation therapy and **ONE** of the following:
      - 1) As a single agent
      - 2) In combination with carboplatin and fluorouracil
      - 3) In combination with cisplatin with or without fluorouracil, docetaxel or paclitaxel
    - ii. If nasopharyngeal site- concurrent radiation and carboplatin
2. Metastatic colorectal cancer (CRC)
  - a. Patient must have **ONE** of the following:
    - i. *KRAS/NRAS* wild-type gene expression as determined by FDA-approved tests **AND ONE** of the following:
      - 1) Used as a single agent: patient has failed oxaliplatin- and irinotecan-based chemotherapy or is intolerant to irinotecan
      - 2) First-line treatment: used in combination with FOLFIRI
      - 3) Used in combination with irinotecan: patients is refractory to irinotecan-based chemotherapy
    - ii. BRAF V600E mutation as detected by an FDA-approved test
      - 1) Used in combination with Braftovi (encorafenib)
      - 2) Patient must **NOT** have wild-type BRAF CRC
      - 3) **NOT** used as first-line therapy
3. Locally advanced or metastatic colorectal cancer (CRC)
  - a. Used in combination with Krazati (adagrasib)
  - b. Patient has received prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy
  - c. Presence of KRAS G12C mutation as determined by an FDA-approved test

4. Metastases of squamous cell skin cancer
5. Metastases of penile cancer
6. Non-small cell lung cancer (NSCLC)
  - a. EGFR mutation
  - b. Progressed after EGFR tyrosine kinase inhibitor therapy
  - c. Used in combination with Gilotrif (afatinib)

**AND** the following for **ALL** indications:

- a. Prescriber agrees to monitor serum electrolytes, magnesium, potassium, calcium levels, and serious infusion reactions.

## Prior - Approval Limits

**Duration** 12 months

---

## Prior – Approval *Renewal* Requirements

**Age** 18 years of age or older

### Diagnoses

Patient must have **ONE** of the following:

1. Squamous cell carcinoma of the head and neck
  - a. Stage III
    - i. If non-nasopharyngeal site- concurrent radiation therapy
  - b. Stage IV
    - i. If non-nasopharyngeal site- concurrent radiation therapy and **ONE** of the following:
      - 1) As a single agent
      - 2) In combination with carboplatin and fluorouracil
      - 3) In combination with cisplatin with or without fluorouracil, docetaxel or paclitaxel
    - ii. If nasopharyngeal site- concurrent radiation and carboplatin
2. Metastatic colorectal cancer (CRC) **AND ONE** of the following:
  - a. Used as a single agent
  - b. Used in combination with FOLFIRI
  - c. Used in combination with irinotecan
  - d. Used in combination with Braftovi encorafenib
3. Locally advanced or metastatic colorectal cancer (CRC) **AND** the following:
  - a. Used in combination with Krazati (adagrasib)
4. Metastases of squamous cell skin cancer



**BlueCross  
BlueShield**

Federal Employee Program.

**ERBITUX  
(cetuximab)**

5. Metastases of penile cancer
6. Non-small cell lung cancer (NSCLC)
  - a. Used in combination with Gilotrif (afatinib)

**AND ALL** of the following for **ALL** indications:

- a. **NO** disease progression or unacceptable toxicity
- b. Prescriber agrees to monitor serum electrolytes, magnesium, potassium, and calcium levels

## **Prior – Approval *Renewal* Limits**

Same as above