

MIEBO
(perfluorohexyloctane ophthalmic solution)

Pre - PA Allowance

None

Prior-Approval Requirements

Age 18 years of age or older

Diagnosis

Patient must have the following:

1. Chronic dry eye
 - a. Patient has been evaluated by an optometrist, ophthalmologist, or a physician specializing in the treatment of the patient's condition
 - b. Prescriber has determined that patient's condition is likely due to meibomian gland dysfunction
 - c. Patient has a Tear Break Up Time (TBUT) indicative of abnormal tear film
 - d. Inadequate treatment response, intolerance, or contraindication to **ONE** product from **EACH** of the following categories of dry eye treatment:
 - i. Cellulose or polyol containing artificial tears (e.g., active ingredients such as: hydroxyethyl cellulose, methylcellulose, Dextran 70, Glycerin, povidone, etc.)
 - ii. Lipid-containing artificial tears (e.g., active ingredients such as: mineral oil, castor oil, flaxseed oil, etc.)
 - iii. Legend ophthalmic for the treatment of dry eyes (see Appendix 1)
 - e. **NO** dual therapy with another legend ophthalmic for the treatment of dry eyes (see Appendix 1)

Prior - Approval Limits

Quantity 3 bottles (9ml) every 90 days

Duration 12 months

Prior – Approval *Renewal* Requirements

Age 18 years of age or older

MIEBO
(perfluorohexyloctane ophthalmic solution)

Diagnosis

Patient must have the following:

1. Chronic dry eye
 - a. Patient has had an improvement in symptoms
 - b. **NO** dual therapy with another legend ophthalmic for the treatment of dry eyes (see Appendix 1)

Prior - Approval *Renewal* Limits

Same as above

Appendix 1 - List of Legend Ophthalmic Medications for Dry Eye

Generic Name	Brand Name
cyclosporine	Cequa
cyclosporine	Restasis
cyclosporine	Vevye
lifitegrast	Xiidra
loteprednol	Eysuvis
perfluorohexyloctane	Miebo
varenicline	Tyrvaya

*Verkazia is not approved for dry eye