

Specialty Guideline Management

Cinqair

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Cinqair	reslizumab

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications¹

Cinqair is indicated for the add-on maintenance treatment of patients with severe asthma aged 18 years and older with an eosinophilic phenotype.

Limitations of Use

- Not for treatment of other eosinophilic conditions
- Not for the relief of acute bronchospasm or status asthmaticus

All other indications are considered experimental/investigational and not medically necessary.

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

Initial requests

- Chart notes or medical record documentation showing pre-treatment blood eosinophil count, or dependence on systemic corticosteroids, if applicable.
- Chart notes, medical record documentation, or claims history supporting previous medications tried including drug, dose, frequency, and duration.

Continuation requests

Chart notes or medical record documentation supporting improvement in asthma control.

Prescriber Specialties

This medication must be prescribed by or in consultation with an allergist/immunologist or pulmonologist.

Coverage Criteria¹⁻⁵

Authorization of 6 months may be granted for adult members who have previously received a biologic drug (e.g., Dupixent, Nucala) indicated for asthma in the past year.

Authorization of 6 months may be granted for treatment of severe asthma when all of the following criteria are met:

- Member is 18 years of age or older.
- Member meets either of the following:
 - Member has a baseline blood eosinophil count of at least 400 cells per microliter.
 - Member is dependent on systemic corticosteroids.
- Member has uncontrolled asthma as demonstrated by experiencing at least one of the following within the past year:
 - Two or more asthma exacerbations requiring oral or injectable corticosteroid treatment
 - One or more asthma exacerbation(s) resulting in hospitalization or emergency medical care visit(s)
 - Poor symptom control (frequent symptoms or reliever use, activity limited by asthma, night waking due to asthma)
- Member has inadequate asthma control despite current treatment with both of the following medications at optimized doses:
 - High-dose inhaled corticosteroid
 - Additional controller (i.e., long-acting beta₂-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline)

- Member will continue to use maintenance asthma treatments (i.e., inhaled corticosteroid and additional controller) in combination with the requested medication.

Continuation of Therapy

Authorization of 12 months may be granted for treatment of severe asthma when all of the following criteria are met:

- Member is 18 years of age or older.
- Asthma control has improved on the requested medication as demonstrated by at least one of the following:
 - A reduction in the frequency and/or severity of symptoms and exacerbations
 - A reduction in the daily maintenance oral corticosteroid dose
- Member will continue to use maintenance asthma treatments (i.e., inhaled corticosteroid and additional controller) in combination with the requested medication.

Other

Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

Note: If the member is a current smoker or vaper, they should be counseled on the harmful effects of smoking and vaping on pulmonary conditions and available smoking and vaping cessation options.

References

1. Cinqair [package insert]. West Chester, PA: Teva Respiratory, LLC; June 2020.
2. Castro M, Zangrilli J, Wechsler ME, et al. Reslizumab for inadequately controlled asthma with elevated blood eosinophil counts: results from two multicentre, parallel, double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet Respir Med*. 2015;3(5):355-366.
3. Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention. 2024 update. Available at: https://ginasthma.org/wp-content/uploads/2024/05/GINA-2024-Strategy-Report-24_05_22_WMS.pdf. Accessed March 1, 2025.
4. American Academy of Allergy, Asthma & Immunology (AAAAI) 2020 Virtual Annual Meeting. Available at: <https://annualmeeting.aaaai.org/>. Accessed March 8, 2025.
5. Cloutier MM, Dixon AE, Krishnan JA, et al. Managing asthma in adolescents and adults: 2020 asthma guideline update from the National Asthma Education and Prevention Program. *JAMA*. 2020;324(22):2301-2317.