

# Specialty Guideline Management

## Rystiggo

### Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Rystiggo	rozanolixizumab-noli

### Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

#### FDA-Approved Indications<sup>1</sup>

Rystiggo is indicated for the treatment of generalized myasthenia gravis (gMG) in adult patients who are anti-acetylcholine receptor (AChR) or anti-muscle-specific tyrosine kinase (MuSK) antibody positive.

All other indications are considered experimental/investigational and not medically necessary.

### Documentation

Submission of the following information is necessary to initiate the prior authorization review:

- For initial requests, chart notes, medical records, or claims history documenting:
  - Positive anti-acetylcholine receptor (AChR) or anti-muscle-specific tyrosine kinase (MuSK) antibody test.
  - Myasthenia Gravis Foundation of America (MGFA) clinical classification.

- MG activities of daily living score.
- Previous medications tried, including response to therapy. If therapy is not advisable, documentation of clinical reasons to avoid therapy.
- For continuation requests: Chart notes or medical record documentation supporting positive clinical response.

## Exclusions

The requested medication will not be used in combination with another neonatal Fc receptor blocker (e.g., Vyvgart, Vyvgart Hytrulo) or complement inhibitor (e.g., Soliris, Ultomiris, Zilbrysq).

## Coverage Criteria

### Generalized Myasthenia Gravis (gMG)<sup>1-4</sup>

Authorization of 6 months may be granted for treatment of generalized myasthenia gravis (gMG) when all of the following criteria are met:

- Anti-acetylcholine receptor (AChR) or anti-muscle-specific tyrosine kinase (MuSK) antibody positive.
- Myasthenia Gravis Foundation of America (MGFA) clinical classification II to IV.
- MG activities of daily living (MG-ADL) total score of greater than or equal to 5.
- Meets one of the following:
  - Member has had an inadequate response or intolerable adverse event to at least two immunosuppressive therapies over the course of at least 12 months (e.g., azathioprine, corticosteroids, cyclosporine, methotrexate, mycophenolate, tacrolimus).
  - Member has had an inadequate response or intolerable adverse event to at least one immunosuppressive therapy and intravenous immunoglobulin (IVIG) over the course of at least 12 months.
  - Member has a documented clinical reason to avoid therapy with immunosuppressive agents and IVIG.

## Continuation of Therapy

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization when there is no evidence of unacceptable toxicity or disease progression while on the current regimen and member demonstrates a positive response to therapy (e.g., improvement in MG-ADL score, MG Manual Muscle Test (MMT), MG Composite).

## References

1. Rystiggo [package insert]. Smyrna, GA: UCB, Inc.; June 2024.
2. Sanders D, Wolfe G, Benatar M et al. International consensus guidance for management of myasthenia gravis. *Neurology*. 2021; 96 (3) 114-122.
3. Bril V, Druzd A, Grosskreutz J, et al. Safety and efficacy of rozanolixizumab in patients with generalised myasthenia gravis (MycarinG): a randomised, double-blind, placebo-controlled, adaptive phase 3 study. *Lancet Neurol*. 2023;22(5):383-394.
4. Barnett C, Herbelin L, Dimachkie MM, Barohn RJ. Measuring Clinical Treatment Response in Myasthenia Gravis. *Neurol Clin*. 2018 May;36(2):339-353.