

STEP THERAPY CRITERIA

BRAND NAME
(generic)

(tacrolimus ointment)

Status: CVS Caremark® Criteria

Type: Initial Step Therapy; Post Step Therapy Prior Authorization

POLICY

FDA-APPROVED INDICATIONS

Tacrolimus ointment, both 0.03% and 0.1% for adults, and only 0.03% for children aged 2 to 15 years, is indicated as *second-line therapy* for the short-term and non-continuous chronic treatment of moderate to severe atopic dermatitis in non-immunocompromised adults and children who have failed to respond adequately to other topical prescription treatments for atopic dermatitis, or when those treatments are not advisable.

Tacrolimus ointment is not indicated for children younger than 2 years of age.

Compendial Uses

Psoriasis³ - on the face, genitals, or skin folds⁵

Vitiligo on the head or neck^{3,6,7}

Atopic Dermatitis for patients under 2 years of age (tacrolimus ointment 0.03%)^{3,4}

INITIAL STEP THERAPY

**Include Rx and OTC products unless otherwise stated.*

For Protopic (tacrolimus) 0.1%, the patient must be at least 16 years of age. For Protopic (tacrolimus) 0.03%, there is no age restriction. Additionally, if the patient has filled a prescription for at least a 14-day supply of at least one corticosteroid of medium or higher potency within the past 180 days (see examples in Table 1) under a prescription benefit administered by CVS Caremark, then the requested drug will be paid under that prescription benefit. If the patient does not meet the initial step therapy criteria, then the system will reject with a message indicating that a prior authorization (PA) is required. The prior authorization criteria would then be applied to requests submitted for evaluation to the PA unit.

TABLE 1: EXAMPLES OF TOPICAL CORTICOSTEROIDS FOR TREATMENT OF ATOPIC DERMATITIS ^{2,3,4}

Medium Potency	betamethasone dipropionate lotion, spray 0.05%
	betamethasone valerate crm/lotion 0.1%/foam 0.12%
	clocortolone pivalate crm 0.1%
	desonide lotion, ointment 0.05%
	desoximetasone crm 0.05%
	fluocinolone acetonide crm/oint/kit 0.025%
	flurandrenolide crm/oint/lotion 0.05%
	fluticasone propionate crm/lotion 0.05%/oint 0.005%
	hydrocortisone butyrate cream/lipocream/lotion/oint/soln 0.1%
	hydrocortisone probutate crm 0.1%
	hydrocortisone valerate crm/oint 0.2%
	mometasone furoate crm/lotion/solution 0.1%
	prednicarbate crm/oint 0.1%
	triamcinolone acetonide crm/oint/lotion/kit 0.1%
	triamcinolone acetonide crm/oint/lotion 0.025%
	triamcinolone acetonide ointment 0.05%
High Potency	amcinonide crm/oint/lotion 0.1%

Tacrolimus Ointment Logic with Age, Post PA Policy UDR 04-2024.docx

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark.

©2024 CVS Health and/or its affiliates. All rights reserved. 106-58428A 021423

	betamethasone dipropionate crm/oint 0.05%
	betamethasone dipropionate augmented crm/lotion 0.05%
	betamethasone valerate oint 0.1%
	desoximetasone crm/oint/spray 0.25%/gel/oint 0.05%
	diflorasone diacetate crm (emollient base) 0.05% diflorasone cream 0.05%
	halcinonide crm/oint 0.1%
	fluocinonide crm/emulsified cream/oint/gel/soln 0.05%
	mometasone furoate oint 0.1%
	triamcinolone acetonide crm/oint 0.5%
	triamcinolone acetonide aerosol soln 0.147 mg/g
Very High Potency	betamethasone dipropionate augmented oint/gel 0.05%
	clobetasol propionate crm/oint/foam/shampoo/gel/lotion/soln/spray 0.05%/cream 0.025%
	diflorasone diacetate oint 0.05%
	flurandrenolide tape 4mcg/cm ²
	halobetasol propionate crm/oint/lotion/kit 0.05%
	fluocinonide crm 0.1%

COVERAGE CRITERIA

Atopic Dermatitis (Eczema)

Authorization may be granted when the requested drug is being prescribed for the short-term and non-continuous chronic treatment of moderate to severe atopic dermatitis (eczema) when ONE of the following criteria are met:

- The request is for tacrolimus 0.03% ointment and ONE of the following criteria is met:
 - The patient is less than 2 years of age
 - The requested drug will be used on sensitive skin areas (e.g., face, genitals, or skin folds)
 - The patient has experienced an inadequate treatment response, intolerance, or contraindication to at least ONE first line therapy agent (e.g., medium or higher potency topical corticosteroid)
- The request is for tacrolimus 0.1% ointment and the following criteria is met:
 - The patient is 16 years of age or older and ONE of the following criteria is met:
 - The requested drug will be used on sensitive skin areas (e.g., face, genitals, or skin folds)
 - The patient has experienced an inadequate treatment response, intolerance, or contraindication to at least one first line therapy agent (e.g., medium or higher potency topical corticosteroid)

Psoriasis

Authorization may be granted when the requested drug is being prescribed for psoriasis on the face, genitals, or skin folds when ONE of the following criteria are met:

- The request is for tacrolimus 0.03% ointment
- The request is for tacrolimus 0.1% ointment and the following criteria is met:
 - The patient is 16 years of age or older

Vitiligo

Authorization may be granted when the requested drug is being prescribed for vitiligo on the head or neck when ONE of the following criteria are met:

- The request is for tacrolimus 0.03% ointment
- The request is for tacrolimus 0.1% ointment and the following criteria is met:
 - The patient is 16 years of age or older

CONTINUATION OF THERAPY

Atopic Dermatitis

Authorization may be granted when the requested drug is being prescribed for the short-term and non-continuous chronic treatment of moderate to severe atopic dermatitis (eczema) when the following criteria is met:

- The patient has achieved or maintained a positive clinical response as evidenced by improvement [e.g., improvement in or resolution of any of the following signs and symptoms: erythema (redness), edema (swelling),

xerosis (dry skin), erosions, excoriations (evidence of scratching), oozing and crusting, lichenification (epidermal thickening), OR pruritus (itching)] and ONE of the following criteria is met:

- The request is for tacrolimus 0.03% ointment
- The request is for tacrolimus 0.1% ointment and the following criteria is met:
 - The patient is 16 years of age or older

Psoriasis

Authorization may be granted when the requested drug is being prescribed for psoriasis on the face, genitals, or skin folds when the following criteria is met:

- The patient has achieved or maintained a positive clinical response as evidenced by improvement (e.g., clear, or almost clear outcome, patient satisfaction, etc.) and ONE of the following criteria is met:
 - The request is for tacrolimus 0.03% ointment
 - The request is for tacrolimus 0.1% ointment and the following criteria is met:
 - The patient is 16 years of age or older

Vitiligo

Authorization may be granted when the requested drug is being prescribed for vitiligo on the head or neck when the following criteria is met:

- The patient has achieved or maintained a positive clinical response as evidenced by improvement (e.g., meaningful repigmentation) and ONE of the following criteria is met:
 - The request is for tacrolimus 0.03% ointment
 - The request is for tacrolimus 0.1% ointment and the following criteria is met:
 - The patient is 16 years of age or older

DURATION OF APPROVAL (DOA)

- 177-F:
 - 2 years of age and older: Initial therapy DOA: 3 months; Continuation of therapy DOA: 12 months
 - Less than 2 years of age: DOA: 3 months
- 1254-F:
 - 2 years of age and older: Initial therapy DOA: 3 months; Continuation of therapy DOA: 36 months
 - Less than 2 years of age: DOA: 3 months

REFERENCES

1. Tacrolimus Ointment [package insert]. Melville, NY: Fougera PHARMACEUTICALS INC; November 2023.
2. Lexicomp Online, AHFS DI (Adult and Pediatric) Online. Waltham, MA: UpToDate, Inc.; 2024. <https://online.lexi.com>. Accessed February 13, 2024.
3. Micromedex® (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: <https://www.micromedexsolutions.com/> (cited: 02/13/2024).
4. Eichenfield L, Tom W, et al. Guidelines of Care for the Management of Atopic Dermatitis. Section 2. Management and Treatment of Atopic Dermatitis with Topical Therapies. *J Am Acad Dermatol*. 2014;71:116-32.
5. Elmetts CA, Korman NJ, Prater EF, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures. *J Am Acad Dermatol*. 2021 Feb;84(2):432-470.
6. Kubelis-López DE, Zapata-Salazar NA, et al. Updates and new medical treatments for vitiligo (Review). *Exp Ther Med*. 2021;22(2):797.
7. Eleftheriadou V, Atkar R, et al. British Association of Dermatologists guidelines for the management of people with vitiligo 2021. *The British Journal of Dermatology*. 2021;186(1):18-29.
8. Eichenfield LF, Tom WL, et. al. Guidelines of Care for the Management of Atopic Dermatitis: Section 1. Diagnosis and Assessment of Atopic Dermatitis. *J Am Acad Dermatol* 2014; 70:338-51.
9. Sidbury RS, Alikhan A, Berovitch L, et al. Guidelines of care for the management of atopic dermatitis in adults with topical therapies. *J Am Acad Dermatol*. 2023; 89(1): e1-e20.