

QUANTITY LIMIT CRITERIA

DRUG CLASS	INSOMNIA AGENTS**
BRAND NAME (generic)	
	AMBIEN (zolpidem)
	AMBIEN CR (zolpidem extended-release)
	DORAL (quazepam)
	(estazolam)
	(flurazepam)
	HALCION (triazolam)
	LUNESTA (eszopiclone)
	RESTORIL (temazepam)
	ROZEREM (ramelteon)
	(zaleplon)
	(zolpidem tartrate capsules)

Status: CVS Caremark® Caremark Criteria

Type: Quantity Limit

***Edluar, zolpidem tartrate sublingual tablets (generic Intermezzo), ZolpiMist, Belsomra, Dayvigo, and Quviviq are not included in these criteria. Refer to Insomnia (zolpidem sublingual, oral spray) or Insomnia (Belsomra, Dayvigo, Quviviq) Prior Authorization criteria.*

POLICY

FDA-APPROVED INDICATIONS

Ambien

Insomnia Limit Policy 01-2024.docx

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark.

©2024 CVS Health and/or its affiliates. All rights reserved. 106-58428A 021423

Ambien (zolpidem tartrate) is indicated for the short-term treatment of insomnia characterized by difficulties with sleep initiation. Ambien has been shown to decrease sleep latency for up to 35 days in controlled clinical studies. The clinical trials performed in support of efficacy were 4-5 weeks in duration with the final formal assessments of sleep latency performed at the end of treatment.

Ambien CR

Ambien CR (zolpidem tartrate extended-release tablets) is indicated for the short-term treatment of insomnia characterized by difficulties with sleep onset and/or sleep maintenance (as measured by wake time after sleep onset). The clinical trials performed in support of efficacy were up to 3 weeks (using polysomnography measurement up to 2 weeks in both adult and elderly patients) and 24 weeks (using patient-reported assessment in adult patients only) in duration.

Doral

Doral is indicated for the treatment of insomnia characterized by difficulty in falling asleep, frequent nocturnal awakenings, and/or early morning awakenings. The effectiveness of Doral has been established in placebo-controlled clinical studies of 5 nights duration in acute and chronic insomnia. The sustained effectiveness of Doral has been established in chronic insomnia in a sleep lab (polysomnographic) study of 28 nights duration. Because insomnia is often transient and intermittent, the prolonged administration of Doral Tablets is generally not necessary or recommended. Since insomnia may be a symptom of several other disorders, the possibility that the complaint may be related to a condition for which there is a more specific treatment should be considered.

Estazolam

Estazolam is indicated for the short-term management of insomnia characterized by difficulty in falling asleep, frequent nocturnal awakenings, and/or early morning awakenings. Both outpatient studies and a sleep laboratory study have shown that estazolam administered at bedtime improved sleep induction and sleep maintenance. Because insomnia is often transient and intermittent, the prolonged administration of estazolam is generally neither necessary nor recommended. Since insomnia may be a symptom of several other disorders, the possibility that the complaint may be related to a condition for which there is a more specific treatment should be considered. There is evidence to support the ability of estazolam to enhance the duration and quality of sleep for intervals up to 12 weeks.

Flurazepam

Flurazepam hydrochloride capsules are indicated for the treatment of insomnia characterized by difficulty in falling asleep, frequent nocturnal awakenings, and/or early morning awakenings. Since insomnia is often transient and intermittent, short-term use is usually sufficient. Prolonged use of hypnotics is usually not indicated and should only be undertaken concomitantly with appropriate evaluation of the patient.

Halcion

Halcion is indicated for the short-term treatment of insomnia (generally 7 to 10 days) in adults.

Lunesta

Lunesta (eszopiclone) is indicated for the treatment of insomnia. In controlled outpatient and sleep laboratory studies, Lunesta administered at bedtime decreased sleep latency and improved sleep maintenance. The clinical trials performed in support of efficacy were up to 6 months in duration. The final formal assessments of sleep latency and maintenance were performed at 4 weeks in the 6-week study (adults only), at the end of both 2-week studies (elderly only) and at the end of the 6-month study (adults only).

Restoril

Restoril (temazepam) is indicated for the short-term treatment of insomnia (generally 7 to 10 days). For patients with short-term insomnia, instructions in the prescription should indicate that Restoril (temazepam) should be used for short periods of time (7 to 10 days). The clinical trials performed in support of efficacy were 2 weeks in duration with the final formal assessment of sleep latency performed at the end of treatment.

Rozerem

Rozerem is indicated for the treatment of insomnia characterized by difficulty with sleep onset.

The clinical trials performed in support of efficacy were up to six months in duration. The final formal assessments of sleep latency were performed after two days of treatment during the crossover study (elderly only), at five weeks in the six week studies (adults and elderly), and at the end of the six month study (adults and elderly).

Zaleplon

Zaleplon capsules are indicated for the short-term treatment of insomnia. Zaleplon capsules have been shown to decrease the time to sleep onset for up to 30 days in controlled clinical studies. They have not been shown to increase total sleep time or decrease the number of awakenings.

The clinical trials performed in support of efficacy ranged from a single night to 5 weeks in duration. The final formal assessments of sleep latency were performed at the end of treatment.

Zolpidem Tartrate Capsules

Zolpidem Tartrate Capsules are indicated for the short-term treatment of transient insomnia characterized by difficulties with sleep initiation in adults younger than 65 years of age.

INITIAL LIMIT QUANTITY		
Limits should accumulate across all drugs and strengths up to highest quantity listed depending on the order the claims are processed.		
Drug	1 Month Limit*	3 Month Limit*
Ambien all strengths (zolpidem)	15 tablets / 25 days	45 tablets / 75 days
Ambien CR all strengths (zolpidem extended-release)	15 tablets / 25 days	45 tablets / 75 days
Doral all strengths (quazepam)	15 tablets / 25 days	45 tablets / 75 days
estazolam all strengths	15 tablets / 25 days	45 tablets / 75 days
flurazepam all strengths	15 capsules / 25 days	45 capsules / 75 days
Halcion all strengths (triazolam)	10 tablets / 25 days	30 tablets / 75 days
Lunesta all strengths (eszopiclone)	15 tablets / 25 days	45 tablets / 75 days
Restoril all strengths (temazepam)	15 capsules / 25 days	45 capsules / 75 days
Rozerem all strengths (ramelteon)	15 tablets / 25 days	45 tablets / 75 days
zaleplon all strengths	15 capsules / 25 days	45 capsules / 75 days
zolpidem tartrate capsules all strengths	15 capsules / 25 days	45 capsules / 75 days
<i>*The duration of 25 days is used for a 30-day fill period and 75 days is used for a 90-day fill period to allow time for refill processing.</i>		

REFERENCES

1. Ambien [package insert]. Bridgewater, NJ: sanofi-aventis U.S. LLC; February 2022.
2. Ambien CR [package insert]. Bridgewater, NJ: sanofi-aventis U.S. LLC; February 2022.
3. Doral [package insert]. Atlanta, GA: Galt Pharmaceuticals, LLC.; January 2023.
4. Estazolam [package insert]. Raleigh, NC: Mayne Pharma; November 2022.
5. Flurazepam Hydrochloride [package insert]. Congers, NY: Chartwell RX, LLC; September 2023.
6. Halcion [package insert]. New York, NY: Pharmacia & Upjohn Co; January 2023.
7. Lunesta [package insert]. Wixom, MI: Waylis Therapeutics LLC; July 2023.

Insomnia Limit Policy 01-2024.docx

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark.

©2024 CVS Health and/or its affiliates. All rights reserved. 106-58428A 021423

8. Restoril [package insert]. Webster Groves, MO: SpecGx LLC; January 2023.
9. Rozerem [package insert]. Lexington, MA: Takeda Pharmaceuticals America, Inc.; November 2021.
10. Zaleplon [package insert]. East Windsor, NJ: Aurobindo Pharma USA, Inc.; October 2021.
11. Zolpidem Tartrate Capsules [package insert]. Morristown, NJ: Almatica Pharma LLC; May 2023.
12. Lexicomp Online, AHFS DI (Adult and Pediatric) Online. Waltham, MA: UpToDate, Inc.; 2023. <https://online.lexi.com>. Accessed November 28, 2023.
13. Micromedex (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: <https://www.micromedexsolutions.com/> (cited: 11/28/2023).
14. Sateia MJ, Buysse DJ, Krystal AD, et al. Clinical practice guideline for the pharmacologic treatment of chronic insomnia in adults: An American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2017;13(2):307-349.
15. American Academy of Sleep Medicine. International Classification of Sleep Disorders, 3rd edition, text revision. American Academy of Sleep Medicine, 2023.
16. Schutte-Rodin S, Broch L, Buysse D, et al. Clinical guideline for the evaluation and management of chronic insomnia in adults. *J Clin Sleep Med*. 2008; 4(5):487-504.
17. Crescenzo FD, D'Alo GL, Ostinelli EG, et al. Comparative effects of pharmacological interventions for the acute and long-term management of insomnia disorder in adults: a systematic review and network meta-analysis. *Lancet*. 2022;400:170-184.