

# Specialty Guideline Management L-glutamine powder-Endari

### **Products Referenced by this Document**

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over the counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Endari	L-glutamine oral powder

# Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

#### **FDA-approved Indication**

Endari is indicated to reduce the acute complications of sickle cell disease in adult and pediatric patients 5 years of age and older.

All other indications are considered experimental/investigational and not medically necessary.

## **Prescriber Specialties**

This medication must be prescribed by or in consultation with a hematologist or specialist in sickle cell disease.

L-glutamine powder-Endari SGM 2208-A P2024\_R.docx

© 2024 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

## **Coverage Criteria**

#### Sickle cell disease, to reduce acute complications

Authorization of 12 months may be granted for use in reducing the acute complications of sickle cell disease in members 5 years of age or older when either of the following criteria is met:

- Member has sickle hemoglobin C (HbSC), sickle β<sup>+</sup>-thalassemia (HbSβ<sup>+</sup>), or other genotypic variants of sickle cell disease (e.g., HbS-O Arab, HbS-Lepore).
- Member has homozygous hemoglobin S (HbSS) or sickle β<sup>0</sup>-thalassemia (HbSβ<sup>0</sup>) genotype AND meets any of the following:
  - Has experienced, at any time in the past, an inadequate response or intolerance to a trial of hydroxyurea
  - Has a contraindication to hydroxyurea
  - Will be using the requested medication with concurrent hydroxyurea therapy

# **Continuation of Therapy**

#### Sickle cell disease, to reduce acute complications

Authorization of 12 months may be granted for continued treatment when the member has experienced a reduction in acute complications of sickle cell disease (e.g., reduction in the number of painful vaso-occlusive episodes, acute chest syndrome episodes, fever, occurrences of priapism, splenic sequestration) since initiating therapy with the requested medication.

### References

- 1. Endari [package insert]. Torrance, CA: Emmaus Medical, Inc; October 2020.
- 2. L-glutamine [package insert]. East Windsor, NJ: Novitium Pharma LLC; July 2024.
- Niihara Y, Miller ST, Kanter J, et al. A phase 3 trial of l-glutamine in sickle cell disease. N Engl J Med. 2018;379(3):226-235.

© 2024 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.