

Reference number(s)
2248-A

Specialty Guideline Management Dysport

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Dysport	abobotulinumtoxin A

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications

- Treatment of cervical dystonia in adults
- Treatment of spasticity in patients 2 years of age and older

Compendial Uses

- Blepharospasm
- Hemifacial spasm
- Chronic anal fissures
- Excessive salivation
- Primary axillary hyperhidrosis

All other indications are considered experimental/investigational and not medically necessary.

Prescriber Specialties

The medication must be prescribed by or in consultation with a provider specialized in the treating the member's condition.

Exclusions

Coverage will not be provided for cosmetic use.

Coverage Criteria

Cervical dystonia

Authorization of 12 months may be granted for treatment of adults with cervical dystonia (e.g., torticollis) when all of the following are met:

- Member is 18 years of age or older
- Member has abnormal placement of the head with limited range of motion in the neck.

Upper or lower limb spasticity

Authorization of 12 months may be granted for treatment of upper or lower limb spasticity when all of the following are met:

- Member is 2 years of age or older
- Member has a primary diagnosis of upper or lower limb spasticity or as a symptom of a condition (including focal spasticity or equinus gait due to cerebral palsy)

Blepharospasm

Authorization of 12 months may be granted for treatment of blepharospasm, including blepharospasm associated with dystonia and benign essential blepharospasm.

Hemifacial spasm

Authorization of 12 months may be granted for treatment of hemifacial spasm.

Chronic anal fissures

Authorization of 12 months may be granted for treatment of chronic anal fissures when the member has not responded to first-line therapy such as topical calcium channel blockers or topical nitrates.

Dysport SGM 2248-A P2024_R.docx

 $\hbox{@ 2024 CVS Caremark.}$ All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

Excessive salivation

Authorization of 12 months may be granted for treatment of excessive salivation (chronic sialorrhea) when the member has been refractory to pharmacotherapy (e.g., anticholinergics).

Primary axillary hyperhidrosis

Authorization of 12 months may be granted for treatment of primary axillary hyperhidrosis when all of the following criteria are met:

- Significant disruption of professional and/or social life has occurred because of excessive sweating; and
- Topical aluminum chloride or other extra-strength antiperspirants are ineffective or result in a severe rash.

Continuation of Therapy

All members (including new members) requesting authorization for continuation of therapy must meet all requirements in the coverage criteria and be experiencing benefit from therapy.

References

- 1. Dysport [package insert]. Wrexham, UK: Ipsen Biopharm, Ltd.; September 2023.
- 2. DRUGDEX® System (electronic version). Truven Health Analytics, Ann Arbor, MI. Available at http://www.micromedexsolutions.com. Accessed August 15, 2024.
- 3. Lexi-Drugs. Hudson, OH: Lexicomp, 2019. http://online.lexi.com/. Accessed August 15, 2024.
- 4. Simpson DM, Hallett M, Ashman EJ et al. Practice guideline update summary: Botulinum neurotoxin for the treatment of blepharospasm, cervical dystonia, adult spasticity, and headache. Report of the Guideline Development Subcommittee of the American Academy of Neurology. Neurology 2016;86:1818-1826.
- 5. Dashtipour K, Chen JJ, Frei K, et al. Systemic Literature Review of AbobotulinumtoxinA in Clinical Trials for Blepharospasm and Hemifacial Spasm. Tremor Other Hyperkinet Mov (NY). 2015;5:338.
- 6. Lakraj AA, Moghimi N, Jabbari B. Sialorrhea: Anatomy, Pathophysiology and Treatment with Emphasis on the Role of Botulinum Toxins. *Toxins* 2013, 5, 1010-1031
- 7. Glader L, Delsing C, Hughes A et al. Sialorrhea in cerebral palsy. American Academy for Cerebral Palsy and Developmental Medicine Care Pathways. https://www.aacpdm.org/publications/care-pathways/sialorrhea. Accessed August 15, 2024.
- 8. Garuti G, Rao F, Ribuffo V et al. Sialorrhea in patients with ALS: current treatment options. *Degener Neurol Neuromuscul Dis.* 2019; 9: 19–26.