

Reference number(s) 4395-A

Specialty Guideline Management Oxlumo

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Oxlumo	lumasiran

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications¹

Oxlumo is indicated for the treatment of primary hyperoxaluria type 1 (PH1) to lower urinary and plasma oxalate levels in pediatric and adult patients.

All other indications are considered experimental/investigational and not medically necessary.

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

Initial requests:

 Molecular genetic test results demonstrating a mutation in the alanine:glyoxylate aminotransferase (AGXT) gene or liver enzyme analysis results demonstrating absent or significantly reduced alanine:glyoxylate aminotransferase (AGT) activity.

Oxlumo SGM 4395-A P2024.docx

© 2024 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

Continuation requests:

Chart notes or medical records demonstrating a positive response to therapy.

Coverage Criteria

Primary Hyperoxaluria Type 1 (PH1)¹⁻³

Authorization of 12 months may be granted for the treatment of primary hyperoxaluria type 1 (PH1) when all of the following criteria are met:

- Member has a diagnosis of PH1 confirmed by either of the following:
 - Molecular genetic test results demonstrating a mutation in the alanine:glyoxylate aminotransferase (AGXT) gene.
 - Liver enzyme analysis results demonstrating absent or significantly reduced alanine:glyoxylate aminotransferase (AGT) activity.
- The requested medication will not be used in combination with nedosiran.

Continuation of Therapy

Authorization of 12 months may be granted for members who meet all requirements in the coverage criteria section and demonstrate a positive response to therapy (e.g., decrease or normalization in urinary and/or plasma oxalate levels, improvement in kidney function).

References

- 1. Oxlumo [package insert]. Cambridge, MA: Alnylam Pharmaceuticals, Inc; September 2023.
- 2. Niaudet, P. Primary hyperoxaluria. In: UpToDate, Post, TW (Ed), UpToDate, Waltham, MA, 2024.
- 3. Milliner DS. The primary hyperoxalurias: an algorithm for diagnosis. Am J Nephrol 2005; 25:154.