

Initial Prior Authorization with Quantity Limit Foundayo Weight Loss Management

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Foundayo	orforglipron

Indications

FDA-approved Indications

Foundayo is indicated in combination with a reduced-calorie diet and increased physical activity to reduce excess body weight and maintain weight reduction long term in adults with obesity or adults with overweight in the presence of at least one weight-related comorbid condition.

Limitations of Use

Concomitant use with another GLP-1 receptor agonist is not recommended.

Coverage Criteria

Reduction in Excess Body Weight, Maintenance of Weight Reduction Long Term

Authorization may be granted when the requested drug will be used with a reduced-calorie diet AND increased physical activity to reduce excess body weight or maintain weight reduction long term in an adult when ALL of the following criteria are met:

- The patient has participated in a comprehensive weight management program that encourages behavioral modification, reduced-calorie diet, AND increased physical activity with continuing follow-up for at least 6 months prior to using drug therapy.
- The patient meets ONE of the following:
 - The patient has a baseline body mass index (BMI) greater than or equal to 30 kg/m². [NOTE: If the patient is transitioning from another drug therapy for weight loss, please consider their baseline BMI at the start of any drug therapy.] [ACTION REQUIRED: Documentation is required for approval.]
 - The patient has a baseline BMI greater than or equal to 27 kg/m². [NOTE: If the patient is transitioning from another drug therapy for weight loss, please consider their baseline BMI at the start of any drug therapy.] [ACTION REQUIRED: Documentation is required for approval.] In addition, the following criteria is met:
 - The patient has at least ONE weight-related comorbid condition (e.g., hypertension, type 2 diabetes mellitus, dyslipidemia). [NOTE: If the patient is transitioning from another drug therapy for weight loss, please consider their weight-related comorbid condition(s) at the start of any drug therapy.] [ACTION REQUIRED: Documentation is required for approval.]

Continuation of Therapy

Reduction in Excess Body Weight, Maintenance of Weight Reduction Long Term

Authorization may be granted when the requested drug will be used with a reduced-calorie diet AND increased physical activity to reduce excess body weight or maintain weight reduction long term in an adult when ALL of the following criteria are met:

- The patient has completed at least 3 months of therapy with the requested drug at a stable maintenance dose.

Reference number(s)
7433-C

- The patient has lost at least 5 percent of baseline body weight OR the patient has continued to maintain their initial 5 percent weight loss. [ACTION REQUIRED: Documentation is required for approval.]

Quantity Limits Apply

The duration of 25 days is used for a 30-day fill period and 75 days is used for a 90-day fill period to allow time for refill processing.

Drug	Dosage	1 Month Limit	3 Month Limit
Foundayo (orforglipron)	0.8 mg, 2.5 mg, 5.5 mg, 9 mg, 14.5 mg, 17.2 mg	30 tablets / 25 days	90 tablets / 75 days

Duration of Approval (DOA)

- 7433-C:
 - Reduction in excess body weight, maintenance of weight reduction long term: Initial therapy DOA: 8 months; Continuation of therapy DOA: 12 months

References

1. Foundayo [package insert]. Indianapolis, IN: Lilly USA, LLC; April 2026.
2. Lexicomp Online, AHFS DI (Adult and Pediatric) Online. Waltham, MA: UpToDate, Inc.; 2026. <https://online.lexi.com>. Accessed April 1, 2026.
3. Micromedex® (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: <https://www.micromedexsolutions.com/> (cited: 04/01/2026).
4. Jensen MD, Ryan DH, Apovian DM, et al. 2013 AHA/ACC/TOS Guideline for the Management of Overweight and Obesity in Adults A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Obesity Society. *Circulation*. 2014;129(suppl 2):S102-S138.
5. Apovian CM, Aronne LJ, Bessesen DH, et al. Pharmacological Management of Obesity: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2015;100(2):342–362.